



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

deals with the alleged phosphorescence in flowering plants, and concludes that the cases cited are either counterfeit or due to electrical phenomena (St. Elmo's fire).—WILLIAM CROCKER.

#### MINOR NOTICES

**The nuclei of Protista.**—The name "Protista," applied by HAECKER to the lowest animals and plants, has failed to receive general acceptance, even among zoologists, and the forms are found under both the protozoa and unicellular plants. In any consideration of the phylogeny of the nucleus these forms must be of great interest, because the nuclei of the metazoa and of the higher algae and fungi are too highly differentiated to throw much light upon such a subject as the origin of the nucleus. A paper by HARTMAN<sup>5</sup> deals almost entirely with the nucleus of protozoa and its significance as the forerunner of the nucleus of the metazoa. Botanists working with the nucleus in the lower algae and fungi, and especially with flagellates, cannot afford to overlook this paper.—CHARLES J. CHAMBERLAIN.

**Symbolae Antillanae.**<sup>6</sup>—In continuation of this important work Professor URBAN in cooperation with several eminent specialists has issued the second and third fascicles of the seventh volume. There are included descriptions of approximately 300 new species, several varieties, and a few new combinations. The following new genera are proposed: *Sarcopilea* of the Urticaceae, *Plethadenia* of the Rutaceae, *Hypocoton* of the Euphorbiaceae, *Ottoschulzia* of the Icacinaceae, *Maga* of the Malvaceae, *Poicilopsis* of the Asclepiadaceae, *Tuerckheimocharis* of the Scrophulariaceae, and *Shaferocharis* of the Rubiaceae.—J. M. GREENMAN.

#### NOTES FOR STUDENTS

**Root-tuberles of non-leguminous plants.**—In an extremely long and somewhat obscure article, which is not made any clearer by the vague illustrations accompanying it, PEKLO<sup>7</sup> gives an account of his studies of the organisms in the root-tuberles of *Alnus* and *Myrica*. With respect to the morphology of the organisms the author adds nothing to what is known from earlier accounts, especially the excellent account of SHIBATA with which he agrees in all essential details. PEKLO finds in the cells of the root-swellings of *Alnus* and *Myrica* masses of filaments with more or less radial arrangement and termi-

<sup>5</sup> HARTMAN, MAX, Die Konstitution der Protistenkerne und ihre Bedeutung für die Zellenlehre. 8vo. pp. v+54. figs. 13. Jena: Gustav Fischer. 1911.

<sup>6</sup> URBAN, I., Symbolae Antillanae seu fundamenta florae Indiae Occidentalnis. Vol. VII, fasc. 2, pp. 161-304, 15 June; fasc. 3, pp. 305-432, 1 October. Leipzig: Fratres Borntraeger. 1912.

<sup>7</sup> PEKLO, J., Die pflanzlichen Aktimonykosen. Centralbl. Bakt. II. 27:451-579. 1910.